Journal of Mammalogy, 83(4):1172–1182, 2002

INDEX TO VOLUME 83

AUTHORS

Aguirre, L. F., 775
Aines, J., 467
Albuja V., L., 1049
Álvarez-Valin, F., 1127
Amano, M., 370
Arends, A., 445
Arjo, W. M., 754
Asa, C. S., 99
Bacigalupe, L. D., 145
Barquez, R. M., 824
Batzli, G. O., 728
Bauman, J. E., 99
Beck, T. D. I., 1020
Bedin, E., 665
Begall, S., 153
Bentzen, P., 893
Bird, B. H., 478
Blake, B. H., 536
Bodkin, J., 893
Bogan, M. A., 386
Bonvicino, C. R., 1071
Boutin, S., 716
Bowyer, R. T., 229
Bozinovic, F., 145
Bradley, A. J., 71
Bradley, R. D., 20, 396
Braggio, E., 1071
Braun, J. K., 1
Bronner, G. N., 290
Browning, T. L., 437
Brunner, S., 135
Buegge, J. J., 260
Burda, H., 153, 569
Burke da Silva, K., 546
Carrier, M., 767
Carroll, D. S., 396
Carvalho, B. A., 58
Castleberry, S. B., 1058
Cerqueira, R., 354
Cervantes, F. A., 1077
Cheng, H. C., 785
Chimimba, C. T., 290
Cimon, N. J., 229
Clark, B. K., 590
Clark, B. S., 590
Clark, K. L., 1039
Cluff, H. D., 188
Cornick, L. A., 159
Cramer, M. J., 743
Crichton, E. G., 489
Daan, S., 153
da Silva, J., 546
Davidson, S. M., 526
Defila, C., 245
de Freitas, T. R. O., 843
Degen, A. A., 645
Demarais, S., 884
Demers, C. L., 734
DeYoung, R. W., 884
Díaz, M. M., 824
Domingo-Roura, X., 907
dos Reis, S. F., 125, 333, 345
Duan, E., 1035
Duarte, L. C., 333, 345
Dudley, J. P., 652
Durden, L. A., 1039
Edge, W. D., 947
Edwards, C. W., 20
Eldridge, M. D. B., 437
Engstrom, M. D., 1097
Fellers, G. M., 167
Ferguson, J. W. H., 290
Fleming, T. H., 489
Flores, D. A., 824
Ford, W. M., 1058
Frank, C. L., 1013
Frutos, S. D., 815
Galloway, G. J., 71
Gao, Z., 91
García-Perea, R., 110
Gava, A., 843
Gee, K. L., 884
Geiser, F., 81
Giacometti, M., 245
Glass, G. E., 478
Glennon, M. J., 734
Gonzales, R. A., 884
González, S., 1127
Green, J. S., 501
Grogan, R. G., 1020
Gross, J. E., 218
Gubbins, C., 178
Harlow, H. J., 1020
Hass, C. C., 934
Hayano, A., 370
Heaney, L. R., 421
Henry, M., 767
Heth, G., 569
Hoffmann, F. G., 408
Holdø, R. M., 652
Holmes, W. G., 576
Honeycutt, R. L., 884
Hoofer, S. R., 31, 40, 49
Huntly, N., 991
Ibáñez, C., 1049
Inouye, R. S., 991
Iriarte, A. W., 608
Iriarte-Díaz, J., 145
Ismail, K., 665
Iwasa, M. A., 852
Jameson, R., 893
Jia, Z., 1035
Jiang, Z., 91
Jiang, Z., 1035
Johnson, B. K., 229
Johnson, R., 290
Juarez, K. M., 925
Juste, J., 1049
Kalcounis-Rüppell, M. C., 614
Kam, M., 645
Kaufman, D. W., 280
Kaufman, G. A., 280
Kawamichi, T., 553
Khokhlova, I. S., 645
Kie, J. G., 229
King, T. L., 1058
Klein, S. L., 478
Kneeland, M. C., 218
Knowlton, F. F., 501
Kolosova, I. E., 458
Körtner, G., 81
Krutzsch, P. H., 489
Kurta, A., 585
Kyle, C. J., 1141
Larsen, K. W., 716
Larson, S., 893
Lee, L. L., 785
Lee, T. E., Jr., 49
Lemos, B., 354
Leslie, D. M., Jr., 590
Lessa, E. P., 408
Li, J., 91
Lim, B. K., 1097
Lindzey, F. G., 199
Lohuis, T., 1020
López-Fuster, M. J., 1087
López-Wilchis, R., 1049
Lorenzo, C., 1077
Loveridge, A. J., 599
Luna, L., 834
Ma, J., 91
Macdonald, D. W., 599
Maestripieri, D., 563
Mahan, C., 546
Mahoney, S. P., 957
Maldonado, J. E., 1127
Mano, T., 1026
Mares, M. A., 1
Marinho-Filho, J., 925
Markowitz, H., 159
Matlack, R. S., 280
Matocq, M. D., 866
Mattevi, M. S., 58
McConnell, E. K., 260
McDowell, L. R., 652
McLoughlin, P. D., 188
McMahon, K. L., 71
McMillan, B. R., 280
Messier, F., 188
Michener, G. R., 999
Millar, J. S., 614
Miyazaki, N., 370
Monteiro, L. R., 125, 333, 345
Monteiro-Filho, E. L. A., 125, 507
Moshkin, M. P., 458
Murray, D. L., 707
Murray, S. W., 585
Nelson, R. J., 478
Novikov, E. A., 458
Nunes, A. P., 58

Journal of Mammalogy, 83(4):1172–1182, 2002

INDEX TO VOLUME 83

AUTHORS

Aguirre, L. F., 775
Aines, J., 467
Albuja V., L., 1049
Álvarez-Valin, F., 1127
Amano, M., 370
Arends, A., 445
Arjo, W. M., 754
Asa, C. S., 99
Bacigalupe, L. D., 145
Barquez, R. M., 824
Batzli, G. O., 728
Bauman, J. E., 99
Beck, T. D. I., 1020
Bedin, E., 665
Begall, S., 153
Bentzen, P., 893
Bird, B. H., 478
Blake, B. H., 536
Bodkin, J., 893
Bogan, M. A., 386
Bonvicino, C. R., 1071
Boutin, S., 716
Bowyer, R. T., 229
Bozinovic, F., 145
Bradley, A. J., 71
Bradley, R. D., 20, 396
Braggio, E., 1071
Braun, J. K., 1
Bronner, G. N., 290
Browning, T. L., 437
Brunner, S., 135
Buegge, J. J., 260
Burda, H., 153, 569
Burke da Silva, K., 546
Carrier, M., 767
Carroll, D. S., 396
Carvalho, B. A., 58
Castleberry, S. B., 1058
Cerqueira, R., 354
Cervantes, F. A., 1077
Cheng, H. C., 785
Chimimba, C. T., 290
Cimon, N. J., 229
Clark, B. K., 590
Clark, B. S., 590
Clark, K. L., 1039
Cluff, H. D., 188
Cornick, L. A., 159
Cramer, M. J., 743
Crichton, E. G., 489
Daan, S., 153
da Silva, J., 546
Davidson, S. M., 526
Defila, C., 245
de Freitas, T. R. O., 843
Degen, A. A., 645
Demarais, S., 884
Demers, C. L., 734
DeYoung, R. W., 884
Díaz, M. M., 824
Domingo-Roura, X., 907
dos Reis, S. F., 125, 333, 345
Duan, E., 1035
Duarte, L. C., 333, 345
Dudley, J. P., 652
Durden, L. A., 1039
Edge, W. D., 947
Edwards, C. W., 20
Eldridge, M. D. B., 437
Engstrom, M. D., 1097
Fellers, G. M., 167
Ferguson, J. W. H., 290
Fleming, T. H., 489
Flores, D. A., 824
Ford, W. M., 1058
Frank, C. L., 1013
Frutos, S. D., 815
Galloway, G. J., 71
Gao, Z., 91
García-Perea, R., 110
Gava, A., 843
Gee, K. L., 884
Geiser, F., 81
Giacometti, M., 245
Glass, G. E., 478
Glennon, M. J., 734
Gonzales, R. A., 884
González, S., 1127
Green, J. S., 501
Grogan, R. G., 1020
Gross, J. E., 218
Gubbins, C., 178
Harlow, H. J., 1020
Hass, C. C., 934
Hayano, A., 370
Heaney, L. R., 421
Henry, M., 767
Heth, G., 569
Hoffmann, F. G., 408
Holdø, R. M., 652
Holmes, W. G., 576
Honeycutt, R. L., 884
Hoofer, S. R., 31, 40, 49
Huntly, N., 991
Ibáñez, C., 1049
Inouye, R. S., 991
Iriarte, A. W., 608
Iriarte-Díaz, J., 145
Ismail, K., 665
Iwasa, M. A., 852
Jameson, R., 893
Jia, Z., 1035
Jiang, Z., 91
Jiang, Z., 1035
Johnson, B. K., 229
Johnson, R., 290
Juarez, K. M., 925
Juste, J., 1049
Kalcounis-Rüppell, M. C., 614
Kam, M., 645
Kaufman, D. W., 280
Kaufman, G. A., 280
Kawamichi, T., 553
Khokhlova, I. S., 645
Kie, J. G., 229
King, T. L., 1058
Klein, S. L., 478
Kneeland, M. C., 218
Knowlton, F. F., 501
Kolosova, I. E., 458
Körtner, G., 81
Krutzsch, P. H., 489
Kurta, A., 585
Kyle, C. J., 1141
Larsen, K. W., 716
Larson, S., 893
Lee, L. L., 785
Lee, T. E., Jr., 49
Lemos, B., 354
Leslie, D. M., Jr., 590
Lessa, E. P., 408
Li, J., 91
Lim, B. K., 1097
Lindzey, F. G., 199
Lohuis, T., 1020
López-Fuster, M. J., 1087
López-Wilchis, R., 1049
Lorenzo, C., 1077
Loveridge, A. J., 599
Luna, L., 834
Ma, J., 91
Macdonald, D. W., 599
Maestripieri, D., 563
Mahan, C., 546
Mahoney, S. P., 957
Maldonado, J. E., 1127
Mano, T., 1026
Mares, M. A., 1
Marinho-Filho, J., 925
Markowitz, H., 159
Matlack, R. S., 280
Matocq, M. D., 866
Mattevi, M. S., 58
McConnell, E. K., 260
McDowell, L. R., 652
McLoughlin, P. D., 188
McMahon, K. L., 71
McMillan, B. R., 280
Messier, F., 188
Michener, G. R., 999
Millar, J. S., 614
Miyazaki, N., 370
Monteiro, L. R., 125, 333, 345
Monteiro-Filho, E. L. A., 125, 507
Moshkin, M. P., 458
Murray, D. L., 707
Murray, S. W., 585
Nelson, R. J., 478
Novikov, E. A., 458
Nunes, A. P., 58

Núñez-Garduño, A., 1049
Ochiai, K., 964
O'Corry-Crowe, G. M., 1111
O'Donnell, C. F. J., 794
Oliveira, L. F. B., 58, 207
Olson, T. L., 199
Osipova, O. A., 458
Ostrow, D. G., 991
Ostrowski, S., 665
Overkamp, G. J. F., 153
Pacheco, V., 834
Penna-Firme, V., 1071
Peppers, L. L., 396
Pereira, M. E., 467
Pérez-Hernández, R., 1087
Peters, S. L., 1097
Piaggio, A. J., 386
Pierson, E. D., 167
Pitt, W. C., 501
Pletscher, D. H., 754
Porter, W. F., 734
Ream, R. R., 754
Reed, D. F., 218
Reich, R. M., 218
Reynolds, P. S., 321
Ribble, D. O., 260
Rickart, E. A., 421
Rosas, F. C. W., 507
Ross, T. G., 31
Roze, U., 381
Ruiz, A., 445
Salazar, M., 1087
Sanderson, J., 608
Schaefer, J. A., 957
Scheckter, J. L., 467
Sicuro, F. L., 207
Silcox, M. T., 804
Simmons, N. B., 40
Sinclair, E. H., 973
Singer, F. J., 682
Smith, M. F., 408
Snider, I., 645
Soriano, P. J., 445
Spicer, G. S., 386
Staedler, M., 893
Stafford, B. J., 553
Stallman, E. L., 576
Sterbing, S. J., 516
Steury, T. D., 707
Stewart, K. M., 229
Strobeck, C., 1141
Sunquist, M. E., 608
Surov, A. V., 458
Susaki, K., 964
Suzuki, H., 852
Swilling, W. R., Jr., 252
Tabaranza, B. R., Jr., 421
Takatsuki, S., 91
Teaford, M. F., 804
Telitsina, A. Y., 458
Testa, J. W., 699
Thomas, D. W., 767
Thorington, R. W., Jr., 553
Todrank, J., 569
Toftegaard, C. L., 71
Tsubota, T., 1026
Valdespino, C., 99
Valdez, E. W., 386
Van Den Bussche, R. A., 31, 40, 49, 815
van Jaarsveld, A. S., 290
Vaudry, R., 767
Ventura, J., 1087
Von Zuben, F. J., 333, 345
Wang, G., 947
Wang, W., 91
Wang, Z., 1035
Welch, K. C., Jr., 260
Westlake, R. L., 1111
Westman, W., 81
Wheatley, M., 716
Wilkinson, G. S., 526
Williams, J. B., 665
Willig, M. R., 743
Willing, R., 245
Wirsing, A. J., 707
Wolf, M., 728
Wolff, J. O., 947
Wood, P. B., 1058
Wooten, M. C., 252
Wurtz, A. E., 260
Yates, T. L., 1077
Yunger, J. A., 267
Zeigenfuss, L. C., 682
Zeppelin, T. K., 973
Zschokke, S., 674

SUBJECTS

(New names in *italic* type)

Abrocoma:
  bennetti, 1
  boliviensis, 1
  budini, 1
  cinerea, 1
  famatina, 1
  oblativus, 1
  schistacea, 1
  uspallata, 1
  vaccarum, 1
Abrocomidae, 1
Abundance:
  Blarina hylophaga, 280
  estimation by track tubes, 734
  Rangifer tarandus, 957
Acomys cahirinus, 645
Activity patterns:
  Chalinolobus tuberculatus, 794
  Corynorhinus townsendii ingens, 590
  Peromyscus (2 spp.), 614
  Spalacopus cyanus, 153
  Spermophilus richardsonii, 999
Adrenocortical responses to cold, 458
Aerodynamics:
  of gliding behavior in Petaurista leucogenys, 553
Aethomys, 290
Afforestation:
  habitat fragmentation from, 290
Age estimation:
  Formosan leaf-nosed bats, 785
Aggressive displacement:
  in jackal species in Zimbabwe, 599
Alabama:
  Peromyscus polionotus ammobates, 252
Alabama beach mouse, 252
Alarm calling:
  Tamias striatus, 546
Alaska:
  Enhydra lutris, 893
  Eumetopias jubatus, 973
  Phoca vitulina, 1111
Alberta:
  Spermophilus richardsonii, 999
Alces alces, 699
Allegheny woodrats, 1058
Allelic diversity/fixation:
  Martes (2 spp.), 907
  Odocoileus virginianus, 31
Allometry:
  body size estimation of extinct castorids, 321
  Oryx leucoryx, 665
Allozymes:
  Mexican lagomorphs, 1077
  Petrogale penicillata, 437
Alouatta pigra, 159
Alpine ibex, 245
American marten, 734
Ampullary gland:
  Tadarida brasiliensis mexicana, 489
Andean mountain cat, 110
Andes:
  Anoura latidens, 445
  Sturnira erythromos, 445
  Tadarida brasiliensis, 445
  Thomasomys *onkiro*, 834
Anoura latidens, 445
Antechinus stuartii, 71, 81
Antipredator behavior:
  alarm calling in Tamias striatus, 546
Arabian oryx, 665
Arctic:
  Pusa hispida, 370
  Ursus arctos, 188
Argentina:
  Abrocoma (3 spp.), 1
  Gracilinanus ignitus, 824

Arizona:
Nasua narica, 934
Armadillo, 815
Arthropod parasites:
of small mammals in Mississippi, 1039
Artiodactyla, 665
Atlantic bottlenose dolphin, 178
Australia:
Petrogale penicillata, 437
Baculum, 489
Balantiopteryx:
infusca, 1049
io, 1049
plicata, 1049
Banding:
Myotis sodalis, 585
Bats:
Anoura latidens, 445
Balantiopteryx (3 spp.), 1049
Carollia perspicillata, 516
Chalinolobus tuberculatus, 794
Corynorhinus townsendii, 167
Corynorhinus townsendii ingens, 590
Cynomops (5 spp.), 1097
Hipposideros terasensis, 785
Mormoopidae, 40
Myotis lucifugus, 767
Myotis occultus, 386
Myotis sodalis, 585
Phyllostomidae, 49
Saccopteryx bilineata, 526
Sturnira erythromos, 445
Tadarida brasiliensis, 445
Tadarida brasiliensis mexicana, 489
Bears:
Ursus americanus, 1020
Ursus arctos, 188
Ursus arctos yesoensis, 1026
Beaver:
body size estimation of extinct, 321
Beech marten, 907
Behavior:
Petaurista leucogenys, 553
Tamias striatus, 546
Belize:
Alouatta pigra, 159
Bioenergetics:
Acomys cahirinus, 645
responses to cold, 458
Biogeography:
Bullimus (3 spp.), 421
Neotoma mexicana, 20
Birthing synchrony, 699
Birth interval:
Ursus arctos yesoensis, 1026
Black bears, 1020
Black howler monkey, 159
Blarina:
brevicauda, 734
carolinensis, 1039
hylophaga, 280
Body condition:
and prey selection, 707
Body fat:
Ursus americanus, 1020
Body mass:
Acomys cahirinus, 645
Formosan leaf-nosed bats, 785
Ursus americanus, 1020
Body size:
Balantiopteryx (3 spp.), 1049
estimating for extinct mammals, 321
Rangifer tarandus, 957
Body temperature. *See also* Thermoregulation
Sciurus carolinensis, 467
Bolivia:
Didelphis albiventris, 354
neotropical savanna bat community, 775
Bot parasites:
of small mammals in Mississippi, 1039
Bottleneck, population:
Enhydra lutris, 893
Odocoileus virginianus, 31
Brain, imaging of:
Antechinus stuartii, 71
Brazil:
Cerdocyon thous, 925
Chironectes minimus, 58
Chrysocyon brachyurus, 925
Coendou prehensilis, 1071
Ctenomys minutus, 843
Didelphis (3 spp.), 58
Gracilinanus (2 spp.), 58
Lutreolina crassicaudata, 58
Lycalopex vetulus, 925
Marmosa murina, 58
Marmosops (2 spp.), 58
Micoureus demerarae, 58
Monodelphis spp., 58
Oxymycterus nasutus, 408
Pecari tajacu, 207
peccaries, 207
Philander (2 spp.), 58
Sotalia (2 spp.), 125
Sotalia guianensis, 507
Sphiggurus (3 spp.), 1071
Sus scrofa, 207
Tayassu pecari, 207
Thrichomys apereoides, 333, 345
Thylamys velutinus, 58
Brown bears, 1026
Brush-tailed rock-wallaby, 437
Bullimus:
bagobus, 421
gamay, 421
luzonicus, 421
California:
Corynorhinus townsendii, 167
Enhydra lutris, 893
Marmota flaviventris, 576
Peromyscus (2 spp.), 614
Spermophilus lateralis, 1013
Canada:
Rangifer tarandus, 957
Spermophilus richardsonii, 999
Ursus arctos, 188
Canids:
coyote, 501, 754
crab-eating fox, 925
fennec fox, 99
hoary fox, 925
jackal, 599
maned wolf, 925
swift fox, 199
wolf, 754
Canine distemper virus, in:
Vulpes velox, 199
Canis:
adustus, 599
latrans, 501, 754
lupus, 754
mesomelas, 599
Capra ibex ibex, 245
Capricornis crispus, 964
Captive populations, reproduction in:
Canis latrans, 501
Hexaprotodon liberiensis, 674
Paguma larvata, 1035
Carollia perspicillata, 516
Castor, 321
Castoroides, 321
Cattle, free-ranging, 229
Cebidae, 159
Cerdocyon thous, 925
Cerrado biome, 925
Cervus elaphus nelsoni, 229
Chalinolobus tuberculatus, 794
Chile:
Oncifelis guigna, 608
Spalacopus cyanus, 153
China:
Procapra gutturosa, 91
Chipmunk, 546, 734, 1039
Chironectes minimus, 58
Chiroptera:
Carollia perspicillata, 516
Corynorhinus townsendii, 167
Emballonuridae, 526
Mormoopidae, 40
Myotis occultus, 386
neotropical savanna bat community, 775
Phyllostomidae, 49, 516
Saccopteryx bilineata, 526
Chromosome polymorphism. *See also* Karyotypes
Ctenomys minutus, 843
Chrysocyon brachyurus, 925
Circadian rhythm:
Spalacopus cyanus, 153
Civet, masked palm, 1035
Clethrionomys:
gapperi, 734
glareolus, 459

Cline:
Didelphis albiventris populations, 354
Cloud forests, bats in, 445
Coast mole, 804
Coati, white-nosed, 934
Coexistence:
Peromyscus (2 spp.), 614
wolves and coyotes, 754
Colorado:
Oreamonas americanus, 218
Ursus americanus, 1020
Communication. *See also* Vocalizations
Microtus (2 spp.), 536
Community structure:
of neotropical savanna bat community, 775
Competition:
density-dependent, 957
elk, mule deer, and cattle, 207
Rangifer tarandus, 957
sex ratio in pigtail macaques, 563
Competitive exclusion:
elk, mule deer, and cattle, 207
Contact zone:
Neotoma fuscipes, 866
Copulatory plugs:
Paguma larvata, 1035
Corridors:
for Oncifelis guigna, 608
Corynorhinus:
townsendii, 167
townsendii ingens, 590
Cotton mouse, 1039
Cotton rat, 1039
Coyote:
dietary overlap with wolves, 754
reproduction in captive wild-caught, 501
Crab-eating fox, 925
Cranial morphology. *See* Skull morphology
Crocidura, 290
Cryptic species:
Neotoma mexicana species group, 20
Cryptomys:
anselli, 569
kafuensis, 569
mechowi, 569
Ctenomyidae, 843
Ctenomys minutus, 843
Cururo, 145, 153
Cuscomys ashaninka, 1
Cynomops:
abrasus, 1097
greenhalli, 1097
mexicanus, 1097
paranus, 1097
planirostris, 1097
Cytochrome *b* gene:
Coendou and Sphiggurus, 1071
Dasypus novemcinctus, 815
Eothenomys (2 spp.), 852
Myotis occultus, 386
Neotoma mexicana species group, 20
Oxymycterus (11 spp.), 408
Sigmodon (11 spp.), 396
Cytochrome oxidase II gene:
Myotis occultus, 386
Dall sheep, 682
Dasypus novemcinctus, 815
Dasyuridae, 71, 81
Deer:
Odocoileus hemionus hemionus, 229
Odocoileus virginianus, 31, 884
Ozotoceros bezoarticus, 1127
Demography changes:
Rangifer tarandus, 957
Dendromys mysticalis, 290
Den ecology:
Ursus arctos, 188
Density-dependent competition, 957
Dental microwear, 804
Desert bighorn sheep, 682
Desert habitat:
water influx in Oryx leucoryx, 665
Development:
of vocalizations in Carollia perspicillata, 516
Developmental thermoenergetics:
Antechinus stuartii, 81
Dialect:
Saccopteryx bilineata vocalizations, 526
Didelphidae, 58, 354, 1087
Didelphimorphia, 1087
Didelphis:
albiventris, 58, 354
albiventris imperfecta, 1087
albiventris pernigra, 1087
aurita, 58
marsupialis, 58, 1087
Diet:
dental wear in moles, 804
Eumetopias jubatus, 973
Marmota flaviventris, 576
overlap between wolves and coyotes, 754
Spermophilus lateralis, 1013
sympatric canids in Brazil, 925
Thomomys talpoides, 991
Digestive physiology:
Procapra gutturosa, 91
Dipodomys ordii, 743
Dispersal:
Peromyscus polionotus ammobates, 252
Phoca vitulina, 1111
Distribution:
Capricornis crispus, 964
Gulo gulo, 1141
Nasua narica, 207
Neotoma magister, 1058
Ozotoceros bezoarticus, 1127
temporospatial distribution of elk, mule deer, and cattle, 207
Tursiops truncatus, 178
Diversity. *See also* Genetic variation; Species diversity
Odocoileus virginianus, 31
Dog-faced bats, 1097
Dolphins:
Sotalia (2 spp.), 125
Sotalia guianensis, 507
Tursiops truncatus, 178
Dominance rank:
Macaca nemestrina, 563
Dusky-footed wood rat, 866
Eastern chipmunk, 546, 734, 1039
Eastern gray squirrel, 467, 1039
Eastern harvest mouse, 1039
Eastern woodrat, 1039
Echimyidae, 333, 345
Echolocation:
Balantiopteryx (3 spp.), 1049
Carollia perspicillata, 516
Ecology:
Blarina hylophaga, 280
grassland fragmentation, 290
Myotis sodalis, 585
Oncifelis guigna, 608
plant-mediated interactions between Thomomys taloides and herbivorous insects, 991
rodent species diversity, 743
Spalacopus cyanus, 145
Ursus arctos, 188
Ecomorphology:
peccaries and feral hogs, 207
Ectoparasites:
of small mammals in Mississippi, 1039
Elephant, African, 652
Elk, 229
Elliot's short-tailed shrew, 280
Ellobius talpinus, 459
Emballonuridae, 526, 1049
Endangered species:
Corynorhinus townsendii ingens, 590
Myotis sodalis, 585
Ozotoceros bezoarticus, 1127
Energetics of reproduction:
Acomys cahirinus, 645
Ursus americanus, 1020
Energy expenditure:
in Antechinus stuartii development, 81
Energy intake:
Acomys cahirinus, 645
Enhydra lutris, 893
Eothenomys:
andersoni, 852
smithii, 852
Epiphyseal gap measurements, 785
Erethizon dorsatum, 381
Erethizontidae, 1071

Escape terrain:
Oreamonas americanus, 218
Estrous cycles:
Vulpes zerda, 99
Eumetopias jubatus, 135, 973
Evolution:
Coendou and Sphiggurus, 1071
Cynomops, 1097
Japanese voles, 853
Phoca vitulina, 1111
Feeding intensity:
effect on sex ratio in Hexaprotodon liberiensis, 674
Feeding type:
Procapra gutturosa, 91
Female promiscuity:
Odocoileus virginianus, 884
Fleas:
of small mammals in Mississippi, 1039
Food availability:
Capricornis crispus, 964
Food consumption:
Oryx leucoryx, 665
Food deprivation:
Sciurus carolinensis, 467
Food distribution:
Marmota flaviventris, 576
Food supplementation:
Peromyscus leucopus, 267
Foraging:
Balantiopteryx (3 spp.), 1049
Chalinolobus tuberculatus, 794
Corynorhinus townsendii, 167
Eumetopias jubatus, 973
Marmota flaviventris, 576
Myotis lucifugus, 767
Rangifer tarandus, 957
Thomomys talpoides, 991
Forearm length measurements, 785
Formosan leaf-nosed bat, 785
Fossils:
body size estimation of beavers, 321
Fossorial mammal, 991
Fox:
crab-eating, 925
fennec, 99
hoary, 925
swift, 199
Fragmentation of forests:
Oncifelis guigna, 608
Functional ecology:
Spalacopus cyanus, 145
Gazelle, Mongolian, 91
Gene flow:
Dasypus novemcinctus, 815
Gulo gulo, 1141
Neotoma magister, 1058
Genetic differentiation:
Neotoma magister, 1058
Ozotoceros bezoarticus, 1127
Genetic drift:
Phoca vitulina, 1111
Genetic structure:
of Gulo gulo populations, 1141
Genetic variation:
contact zone Neotoma fuscipes, 866
Dasypus novemcinctus, 815
Enhydra lutris, 893
Eothenomys (2 spp.), 852
Mexican lagomorphs, 1077
Neotoma magister, 1058
Odocoileus virginianus, 31
Phoca vitulina, 1111
Geographical information system:
Oreamonas americanus, 218
Geographic variation:
Didelphis albiventris, 354
Odocoileus virginianus, 31
Ozotoceros bezoarticus, 1127
Phoca vitulina, 1111
Pusa hispida, 370
Sotalia (2 spp.), 125
Thrichomys apereoides, 333, 345
in vocalizations of Saccopteryx bilineata, 526
Geometric morphometrics:
Sotalia (2 spp.), 125
Thrichomys apereoides, 333, 345
Geomyidae, 991
Geophagy:
Loxodonta africana, 652
Gestation:
Acomys cahirinus, 645
Chalinolobus tuberculatus, 794
Myotis lucifugus, 767
Sotalia guianensis, 507
Spermophilus richardsonii, 999
Ursus americanus, 1020
Vulpes zerda, 99
Glans penes:
Neotoma fuscipes, 866
Glaucomys sabrinus, 734
Gliding behavior:
Petaurista leucogenys, 553
Golden-mantled ground squirrels, 1013
Golden mouse, 1039
Gracilinanus:
aceramarcae, 824
agilis, 58, 824
dryas, 824
emiliae, 58, 824
ignitus, 824
longicaudus, 824
marica, 824
microtarsus, 58, 824
Grassland fragmentation, 290
Gray squirrel, 1039
Gray-tailed voles, 947
Grizzly bears
Guigna, 608
Gular gland:
Tadarida brasiliensis mexicana, 489
Gulo gulo, 1141
Gut fill:
Procapra gutturosa, 91
Habitat associations:
in rodents, 743
Habitat fragmentation:
Oncifelis guigna, 608
small-mammal assemblages and, 290
Habitat heterogeneity:
in rodents, 743
Habitat model:
Oreamonas americanus, 218
Tamiasciurus hudsonicus, 716
Habitat partitioning:
jackal species in Zimbabwe, 599
Peromyscus (2 spp.), 614
Habitat quality:
Tamiasciurus hudsonicus, 716
Habitat use:
Corynorhinus townsendii, 167
jackal species in Zimbabwe, 599
Oncifelis guigna, 608
sympatric canids in Brazil, 925
Ursus arctos, 188
Vulpes velox, 199
Hairy-tailed mole, 804
Hamsters:
response to cold, 458
Hantavirus:
in Norway rats, 478
Harbor seal, 1111
Hares, snowshoe, 707
Harvest analysis, 1026
Hawaii:
Petrogale penicillata, 437
Hearing:
Carollia perspicillata, 516
Hemicentetes nigriceps, 804
Heterothermy:
Sciurus carolinensis, 467
Hexaprotodon liberiensis, 674
Hibernaculum:
Corynorhinus townsendii ingens, 590
Spermophilus richardsonii, 999
Hibernation:
Corynorhinus townsendii ingens, 590
Spermophilus lateralis, 1013
Spermophilus richardsonii, 999
Ursus americanus, 1020
Hippopotamus, pgymy, 674
Hipposideridae, 785
Hipposideros terasensis, 785
Hoary fox, 925
Hogs, feral, 207
Home range:
Chalinolobus tuberculatus, 794
Myotis lucifugus, 767
Nasua narica, 934
Peromyscus (2 spp.), 260
sympatric canids in Brazil, 925
Tursiops truncatus, 178
Horn growth:
Capra ibex ibex, 245

Howling monkeys, 159
Hunting, influence on mountain sheep, 682
Hybrid:
  hybrid zone in Ctenomys minutus populations, 843
  sea lion, 135
Hystricognathi, 1071
Idaho:
  Canis latrans, 501
  Thomomys talpoides, 991
Illinois:
  Peromyscus leucopus, 267
Indiana bat, 585
Infanticide:
  Microtus canicaudus, 947
Insects:
  insect herbivore interaction with northern pocket gopher, 991
  parasitic arthropods of small mammals, 1039
Interspecific variability:
  response to cold in murine rodents, 458
Iowa:
  Odocoileus virginianus, 31
Isolation-by-distance, 1111
Jackrabbits, 1077
Japan:
  Capricornis crispus, 964
  Eothenomys (2 spp.), 852
  Petaurista leucogenys, 553
  Phoca vitulina, 1111
  Ursus arctos yesoensis, 1026
Japanese giant flying squirrels, 553
Japanese serow, 964
Juvenile recruitment. *See* Recruitment
Kalahari sand habitat, 652
Kansas:
  Blarina hylophaga, 280
Karyotypes:
  Abrocoma uspallata, 1
  Coendou and Sphiggurus, 1071
  Ctenomys minutus, 843
  marsupials from Brazil, 58
  Petrogale penicillata, 437
Kinship odors:
  Cryptomys (3 spp.), 569
Kodkod, 608
Lactation:
  Chalinolobus tuberculatus, 794
  energy cost in Acomys cahirinus, 645
  energy cost in Antechinus stuartii, 81
  Myotis lucifugus, 767
  Spermophilus richardsonii, 999
  Ursus americanus, 1020
Lagomorphs, genetic variation in Mexican, 1077
Lepus:
  americanus, 707
  californicus, 1077
  callotis, 1077
  flavigularis, 1077
Lice:
  of small mammals in Mississippi, 1039
Licks, mineral, 652
Litter size:
  Ursus arctos yesoensis, 1026
  Vulpes velox, 199
Little brown bats, 386, 767
Livetrapping. *See* Trapping
Locomotion:
  gliding behavior in Petaurista leucogenys, 553
Logistic regression:
  Oreamonas americanus, 218
Lophostoma, 49
Loxodonta africana, 652
Lutreolina crassicaudata, 58
Lycalopex vetulus, 925
Lynchailurus pajeros, 110
Macaques, 563
Magnetic resonance imaging:
  Antechinus stuartii, 71
Major histocompatibility complex:
  Odocoileus virginianus, 31
Maned wolf, 925
Marmosa murina, 58
Marmosops:
  incanus, 58
  parvidens, 58
Marmota flaviventris, 576
Marsupials:
  Antechinus stuartii, 71, 81
  developmental thermoenergetics in, 81
  Didelphis (3 spp.), 58, 1087
  Didelphis albiventris, 354
  Petrogale penicillata, 437
  urinary pheromones, 71
Martes:
  americana, 734
  foina, 907
  martes, 907
Maryland:
  hantavirus in Norway rats, 478
Masked palm civets, 1035
Mastomys natalensis, 290
Maternal investment:
  Antechinus stuartii, 81
  Macaca nemestrina, 563
Maternity cave, 590
Mating:
  Capricornis crispus, 964
  hybrid zone in Ctenomys minutus populations, 843
  Odocoileus virginianus, 884
  Ovis (3 spp.), 682
  Paguma larvata, 1035
  Tadarida brasiliensis mexicana, 489
  Vulpes zerda, 99
Maximum likelihood analysis, 386
Meetings, scientific:
  American Society of Mammalogists, 1158
Meles meles, 907
Metabolic rate:
  Antechinus stuartii, 81
  bats from cloud forests, 445
Mexico:
  Balantiopteryx (3 spp.), 1049
  Lepus (3 spp.), 1077
  Neotoma mexicana, 20
  Romerolagus diazi, 1077
  Sylvilagus floridanus, 1077
Michigan:
  Myotis sodalis, 585
Micoureus demerarae, 58
Microsatellite DNA studies:
  Enhydra lutris, 893
  Gulo gulo, 1141
  Martes (2 spp.), 907
  Neotoma fuscipes, 866
  Neotoma magister, 1058
  Odocoileus virginianus, 884
Microtus:
  canicaudus, 947
  montanus, 536
  ochrogaster, 536
  pinetorum, 1039
Migration:
  Myotis sodalis, 585
  Rangifer tarandus, 957
Minimum convex polygon technique, 260
Mississippi:
  Blarina carolinensis, 1039
  Microtus pinetorum, 1039
  Neotoma floridana, 1039
  Ochrotomys nuttalli, 1039
  Oryzomys palustris, 1039
  Peromyscus (2 spp.), 1039
  Rattus norvegicus, 1039
  Reithrodontomys humulis, 1039
  Sciurus carolinensis, 1039
  Sigmodon hispidus, 1039
  Tamias striatus, 1039
Mites:
  of small mammals in Mississippi, 1039
Mitochondrial DNA:
  Coendou and Sphiggurus, 1071
  Dasypus novemcinctus, 815
  Enhydra lutris, 893
  Eothenomys (2 spp.), 852
  Mormoopidae, 40
  Myotis occultus, 386
  Neotoma fuscipes, 866
  Neotoma mexicana species group, 20
  Petrogale penicillata, 437
  Phoca vitulina, 1111
  Sigmodon (11 spp.), 396
  Tonatia, 49
Mole:
  coast, 804
  hairy-tailed, 804

Molecular clock, 386
Molecular phylogenetics:
  Cynompos (5 spp.), 1097
  Petrogale penicillata, 437
  Sigmodon (11 spp.), 396
  Tonatia, 49
Molecular systematics:
  Neotoma mexicana, 20
  Sigmodon (11 spp.), 396
Mole rats, 569
Molossops:
  neglectus, 1097
  temminckii, 1097
Mongolia:
  Procapra gutturosa, 91
Monodelphis:
  brevicaudata, 58
  dimidiata, 58
  domestica, 58
  kunsi, 58
Monogamy:
  Capricornis crispus, 964
  Peromyscus polionotus ammobates, 252
Montana:
  Canis (2 spp.), 754
Montane voles, 536
Moose, 699
Mormoopidae, 40
Mormoops:
  blainvilli, 40
  megalophylla, 40
Morphological key:
  Oreailurus jacobita, 110
Morphology:
  Abrocoma (9 spp.), 1
  body size estimation of extinct castorids, 321
  Bullimus (3 spp.), 421
  Cynomops (5 spp.), 1097
  Didelphis (3 spp.), 354, 1087
  Gracilinanus ignitus, 824
  Mormoopidae, 40
  Neotoma fuscipes, 866
  Oreailurus jacobita, 110
  Otariidae, 135
  Oxymycterus (11 spp.), 408
  Ozotoceros bezoarticus, 1127
  peccaries and feral hogs, 207
  Pusa hispida, 370
  Sotalia (2 spp.), 125
  Spalacopus cyanus, 145
  Thomasomys onkiro, 834
  Thrichomys apereoides, 333, 345
Morphometrics:
  Cynomops (5 spp.), 1097
  Didelphis (3 spp.), 1087
  Didelphis albiventris, 354
  Gracilinanus ignitus, 824
  Otariidae, 135
  Oxymycterus (11 spp.), 408
  Ozotoceros bezoarticus, 1127
  Pusa hispida, 370
  Sotalia (2 spp.), 135
  Spalacopus cyanus, 145
  Thrichomys apereoides, 333, 345
Mortality:
  Nasua narica, 934
Mountain goat, 218
Mountain sheep, 682
Mouse:
  Alabama beach, 252
  cotton, 1039
  eastern harvest, 1039
  golden, 1039
  spiny, 645
  white-footed, 267, 729, 1039
Mule deer, 229
Muridae, 421
Mus minutoides, 290
Mustelid:
  Gulo gulo, 1141
Myosorex varius, 290
Myotis:
  lucifugus, 386, 767
  occultus, 386
  sodalis, 585
Napaeozapus insignis, 734
Nasua narica, 934
Neonatal mortality:
  Ursus arctos yesoensis, 1026
Neotoma:
  distincta, 20
  ferruginea, 20
  floridana, 1039
  fuscipes, 866
  macrotis, 866
  magister, 1058
  mexicana, 20
  navus, 20
  parvidens, 20
  torquata, 20
  tropicalis, 20
Neotropical savanna, 775
Neotropics:
  Didelphis (3 spp.), 1087
  Didelphis albiventris, 354
Nested-clade analysis, 815
Newfoundland:
  Rangifer tarandus, 957
New Mexico:
  Peromyscus (2 spp.), 260
New York:
  Blarina brevicauda, 734
  Clethrionomys gapperi, 734
  Glaucomys sabrinus, 734
  Martes americana, 734
  Napaeozapus insignis, 734
  Odocoileus virginianus, 31
  Sorex cinereus, 734
  Tamiasciurus hudsonicus, 734
  Tamias striatus, 734
  Zapus hudsonius, 734
New Zealand long-tailed bats, 794
Niche partitioning:
  elk, mule deer, and cattle, 207
  peccaries and feral hogs, 207
  Peromyscus (2 spp.), 614
Noctilionoidea, 40
Nocturnal activity:
  Chalinolobus tuberculatus, 794
North America. *See also* Canada; Mexico; United States
  Enhydra lutris, 893
  Erethizon dorsatum, 381
  Gulo gulo, 1141
  Ursus arctos, 188
Northern pocket gopher, 991
Northwest Territories:
  Ursus arctos, 188
Norway rat, 478, 1039
Nunavut:
  Ursus arctos, 188
Ochrotomys nuttalli, 1039
Octodontidae, 145, 153
Odocoileus:
  hemionus hemionus, 229
  virginianus, 31, 884
Odors:
  Cryptomys (5 spp.), 569
Oklahoma:
  Corynorhinus townsendii ingens, 590
  Odocoileus virginianus, 31
Olfaction:
  marsupial, 71
Oncifelis guigna, 608
Ontogeny:
  Carollia perspicillata, 516
Onychomys leucogaster, 743
Opossum, white-eared, 354
Oreailurus jacobita, 110
Oreamonas americanus, 218
Oregon:
  Microtus canicaudus, 947
Oryx leucoryx, 665
Oryzomys palustris, 1039
Otaria byronia, 135
Otariidae, 135
Otomys irroratus, 290
Ovis:
  canadensis canadensis, 682
  canadensis nelsoni, 682
  dalli, 682
Ovulation:
  Sotalia guianensis, 507
  Vulpes zerda, 99
Oxymycterus:
  amazonicus, 408
  dasytrichus, 408
  delator, 408
  hiska, 408
  josei, 408
  judex, 408
  misionalis, 408
  nasutus, 408
  paramensis, 408
  quaestor, 408
  rufus, 408
Ozark big-eared bat, 590
Ozotoceros:
  bezoarticus, 1127
  bezoarticus arerunguaensis, 1127

bezoarticus celer, 1127
bezoarticus leucogaster, 1127
bezoarticus uruguayensis, 1127
Paguma larvata, 1035
Palaeocastor, 321
Pampas cat, 110
Pampas deer, 1127
Paraguay:
Dasypus novemcinctus, 815
Parascalops breweri, 804
Partitioning. *See* Habitat partitioning; Resource partitioning
Parturition:
Ursus arctos yesoensis, 1026
Parturition site:
Spermophilus richardsonii, 999
Paternity, multiple:
Odocoileus virginianus, 884
Pecari tajacu, 207
Peccary, 207
Perognathus flavescens, 743
Peromyscus:
boylii, 260, 614
californicus, 614
gossypinus, 1039
leucopus, 267, 728, 743, 1039
maniculatus, 743
polionotus ammobates, 252
truei, 260
Peru:
Thomasomys onkiro, 834
Petaurista leucogenys, 553
Petrogale penicillata, 437
Pheromones:
Antechinus stuartii, 71
Philander:
frenata, 58
opossum, 58
Philippines:
Bullimus (3 spp.), 421
Philopatry:
Myotis sodalis, 585
Peromyscus polionotus ammobates, 252
Phoca vitulina, 1111
Phodopus:
campbelli, 459
roborovskii, 459
sungorus, 459
Photoperiodicity:
Sciurus carolinensis, 467
Spalacopus cyanus, 153
Phyllostomidae, 49, 517
Phylobiogeography:
Neotoma mexicana, 20
Phoca vitulina, 1111
Phylogenetics:
Cynomops (5 spp.), 1097
Mormoopidae, 40
Myotis occultus, 386
Neotoma mexicana, 20
Phoca vitulina, 1111
Sigmodon (11 spp.), 396
Tonatia, 49
Phylogeny:
body size estimation of extinct castorids, 321
Cynomops (5 spp.), 1097
Eothenomys (2 spp.), 852
Mormoopidae, 40
Odocoileus virginianus, 31
Phoca vitulina, 1111
Phytophenology:
Capra ibex ibex, 245
Pigtail macaques, 563
Pine marten, 907
Pine vole, 1039
Placental scars:
in Canis latrans, 501
Plant-herbivore interaction, 991
Plant-stress hypothesis, 991
Pocket gopher, northern, 991
Polyunsaturated fatty acids (PUFA), effect on ground squirrels, 1013
Population density:
Capricornis crispus, 964
Microtus canicaudus, 947
Nasua narica, 934
Peromyscus leucopus, 267
red squirrel habitat quality, 716
Population diversity:
estimating using track tubes, 734
Population dynamics:
Blarina hylophaga, 280
Ursus arctos yesoensis, 1026
Population fragmentation:
Gulo gulo, 1141
Population genetics:
Neotoma magister, 1058
Porcupine, 381
Postnatal growth in Formosan leaf-nosed bats, 785
Predation:
alarm calling in Tamias striatus, 546
body condition and prey selection, 707
of Nasua narica, 934
selection for earlier births, 699
Predator swamping, 699
Pregnancy. *See* Gestation
Prey identification:
for Eumetopias jubatus, 973
Prey selection:
body condition and, 707
Primates:
Alouatta pigra, 159
Macaca nemestrina, 563
Principal component analysis, 1097
Procapra gutturosa, 91
Promiscuity, female:
Odocoileus virginianus, 884
Prostate:
Tadarida brasiliensis mexicana, 489
Pteromyinae, 553
Pteronotus:
davyi, 40
gymnonotus, 40
macleayii, 40
parnellii, 40
personatus, 40
quadridens, 40
Pusa hispida, 370
Pygmy hippopotamus, 674
Quill, porcupine, 381
Rabbits, 1077
Radiotelemetry:
Corynorhinus townsendii, 167
Lepus americanus, 707
Myotis sodalis, 585
Nasua narica, 934
Oncifelis guigna, 608
Peromyscus (2 spp.), 260
Spermophilus richardsonii, 999
Tamiasciurus hudsonicus, 707
Ursus arctos, 188
Vulpes velox, 199
Rangifer tarandus, 957
Rattus norvegicus, 478, 1039
Recruitment:
Microtus canicaudus, 947
Ovis (3 spp.), 682
Rangifer tarandus, 957
Reithrodontomys humulis, 1039
Reithrodontomys montanus, 743
Reproduction:
Acomys cahirinus, 645
Canis latrans, 501
Chalinolobus tuberculatus, 794
Hexaprotodon liberiensis, 674
Microtus canicaudus, 947
Myotis lucifugus, 767
Odocoileus virginianus, 884
Paguma larvata, 1035
predation selection for earlier births, 699
Sotalia guianensis, 507
Tadarida brasiliensis mexicana, 489
Ursus americanus, 1020
Ursus arctos yesoensis, 1026
Vulpes velox, 199
Resource availability:
red squirrel population density and, 716
Resource partitioning:
elk, mule deer, and cattle, 229
peccaries and feral hogs, 207
Peromyscus (2 spp.), 614
wolves and coyotes, 754
Restriction site mapping, 1097
Rhabdomys pumilio, 290
Ribosomal DNA sequences:
Mormoopidae, 40
Tonatia, 49
Rice rat, 1039
Richardson's ground squirrels, 999
Ringed seals, 370
Rocky Mountain bighorn sheep, 682

Rodentia:
Abrocoma (9 spp.), 1
Acomys cahirinus, 645
Blarina carolinensis, 1039
Blarina hylophaga, 280
Bullimus (3 spp.), 421
Clethrionomys glareolus, 459
Coendou prehensilis, 1071
Cryptomys (5 spp.), 569
Ctenomys minutus, 843
Ellobius talpinus, 459
habitat associations, 743
Microtus canicaudus, 947
Microtus pinetorum, 1039
Neotoma floridana, 1039
Neotoma fuscipes, 866
Neotoma magister, 1058
Neotoma mexicana, 20
Ochrotomys nuttalli, 1039
Oryzomys palustris, 1039
Peromyscus (2 spp.), 260, 1039
Peromyscus leucopus, 267, 728, 1039
Peromyscus polionotus ammobates, 252
Phodopus (3 spp.), 459
Rattus norvegicus, 478, 1039
Reithrodontomys humulis, 1039
Sciurus carolinensis, 1039
Sigmodon hispidus, 1039
Spalacopus cyanus, 145, 153
Spermophilus lateralis, 1013
Spermophilus richardsonii, 999
Sphiggurus (2 spp.), 1071
Tamias striatus, 564, 1039
Thomasomys onkiro, 834
Thomomys talpoides, 991
Thrichomys apereoides, 333, 345
Romerolagus diazi, 1077
Roost sites:
Chalinolobus tuberculatus, 794
Corynorhinus townsendii, 167
Corynorhinus townsendii ingens, 590
Myotis lucifugus, 767
Myotis sodalis, 585
Saccopteryx bilineata, 526
Sac-winged bats, 526, 1049
Sagebrush steppe:
Vulpes velox, 199
Sampling gap, 345
Saudi Arabia:
Oryx leucoryx, 665
Scapanus orarius, 804
Scat analysis:
Eumetopias jubatus, 973
Sciuridae, 576
Sciurus carolinensis, 467, 1039
Sea lions, Steller, 973
Seals:
harbor, 1111
ringed, 370
Sea otter, 893
Seasonality:
Corynorhinus townsendii ingens, 590
in digestive strategy of Procapra gutturosa, 91
Eumetopias jubatus prey consumption, 973
of heterothermy in Sciurus carolinensis, 467
Oreamnos americanus, 218
of reproductive biology of Tadarida brasiliensis mexicana, 489
of Seoul virus in Norway rats, 478
Seminal vesicle:
Tadarida brasiliensis mexicana, 489
Senescence:
Canis latrans, 501
Sotalia guianensis, 507
Seoul virus in Norway rats, 478
Serow, Japanese, 964
Sex-biased capture:
Peromyscus leucopus, 728
Sex-determining gene of Y chromosome (*Sry*):
Eothenomys (2 spp.), 852
Sex ratio:
Capricornis crispus, 964
Hexaprotodon liberiensis, 674
Macaca nemestrina, 563
Microtus canicaudus, 947
Sexual dimorphism:
Didelphis (3 spp.), 1087
Ozotoceros bezoarticus, 1127
Sexual maturity:
Microtus canicaudus, 947
Ursus arctos yesoensis, 1026
Shrew:
Crocidura, 290
Elliot's short-tailed, 280
southern short-tailed, 1039
Sibling species:
Neotoma mexicana species group, 20
Sigmodon:
alleni, 396
alstoni, 396
arizonae, 396
fulviventer, 396
hirsutus, 396
hispidus, 396, 1039
leucotis, 396
mascotensis, 396
ochrognathus, 396
peruanus, 396
toltecus, 396
Sigmodontinae, 834
Skull morphology:
Didelphis (3 spp.), 1087
Didelphis albiventris, 354
Gracilinanus ignitus, 824
Otariidae, 135
Pecari tajacu, 207
Pusa hispida, 370
Sotalia (2 spp.), 125
Sus scrofa, 207
Tayassu pecari, 207
Thomasomys onkiro, 834
Thrichomys apereoides, 333, 345
Sleep-site use:
Spermophilus richardsonii, 999
Snowshoe hares, 707
Sodium availability:
Loxodonta africana, 652
Sorex cinereus, 734
Sotalia:
fluviatilis, 125
guianensis, 125, 507
South Africa:
grassland fragmentation, 290
South America:
Abrocoma (9 spp.), 1
Anoura latidens, 445
Cerdocyon thous, 925
Chrysocyon brachyurus, 925
Ctenomys minutus, 843
Dasypus novemcinctus, 815
Didelphis (3 spp.), 1087
Didelphis albiventris, 354
Gracilinanus ignitus, 824
Lycalopex vetulus, 925
Lynchailurus pajeros, 110
neotropical savanna bat community, 775
Oncifelis guigna, 608
Oreailurus jacobita, 110
Oxymycterus (11 spp.), 408
Ozotoceros bezoarticus, 1127
Saccopteryx bilineata, 526
Sotalia (2 spp.), 125
Sotalia guianensis, 507
Sturnira erythromos, 445
Tadarida brasiliensis, 445
Thomasomys onkiro, 834
Thrichomys apereoides, 333, 345
South Carolina:
Tursiops truncatus, 178
Southern short-tailed shrew, 1039
Spalacopus cyanus, 145, 153
Speciation:
Bullimus (3 spp.), 421
Oxymycterus (11 spp.), 408
Species diversity:
habitat heterogeneity and, 743
of neotropical savanna bat community, 775
Species odors:
Cryptomys (3 spp.), 569
Spermatozoa:
Paguma larvata, 1035
Tadarida brasiliensis mexicana, 489
Spermophilus:
lateralis, 1013
richardsonii, 999
Sphiggurus:
insidiosus, 1071
melanura, 1071
vilosus, 1071

Spiny mouse, 645
Squirrels:
  eastern gray, 467
  golden-mantled ground, 1013
  Japanese giant flying, 553
  red, 707, 716
  Richardson's ground, 999
Steller sea lions, 973
Stress reactivity:
  to cold in murine rodents, 458
Sturnira erythromos, 445
Subspecies:
  Ozotoceros bezoarticus, 1127
  Pusa hispida, 370
Subterranean rodent:
  Spalacopus cyanus, 145, 153
Survival rates:
  Microtus canicaudus, 947
  Nasua narica, 934
  Ovis (3 spp.), 682
  Rangifer tarandus, 957
  Vulpes velox, 199
Sus scrofa, 207
Swift fox, 199
Switzerland:
  Capra ibex ibex, 245
Sylvilagus floridanus, 1077
Sympatry:
  canids in Brazil, 925
  Didelphis (3 spp.), 1087
  jackal species in Zimbabwe, 599
  Martes (2 spp.), 907
Systematics:
  Abrocoma cinerea species complex, 1
  Coendou and Sphiggurus, 1071
  Cynomops, 1097
  Eothenomys (2 spp.), 852
  Gracilinanus ignitus, 824
  Myotis occultus, 386
  Neotoma mexicana species group, 20
  Oxymycterus (11 spp.), 408
  Sigmodon (11 spp.), 396
  Tonatia, 49
Tadarida:
  brasiliensis, 445
  brasiliensis mexicana, 489
Tallgrass prairie:
  Blarina hylophaga, 280
Tamiasciurus hudsonicus, 707, 716, 734
Tamias striatus, 546, 734, 1039
Taxonomy:
  Abrocoma cinerea species complex, 1
  Coendou and Sphiggurus, 1071
  Cynomops (5 spp.), 1097
  Didelphis (3 spp.), 1087
  Didelphis albiventris, 354
  Gracilinanus ignitus, 824
  Mormoopidae, 40
  Myotis occultus, 386
  Neotoma fuscipes, 866
  Petrogale penicillata, 437
  Tonatia, 49
Tayassuidae, 207
Tayassu pecari, 207
Temperature, effects of:
  adrenocortical and bioenergetic responses to cold in murine rodents, 458
  horn growth in Capra ibex ibex, 245
  physiology of bats from Andean cloud forests, 445
  Seoul virus in Norway rats, 478
  Spermophilus lateralis diet, 1013
Temporospatial distribution:
  elk, mule deer, and cattle, 207
Tennessee:
  Odocoileus virginianus, 31
Tenrecs, 804
Territoriality:
  Capricornis crispus, 964
  Microtus canicaudus, 947
Territory size:
  Capricornis crispus, 964
Testis:
  Sotalia guianensis, 507
  Tadarida brasiliensis mexicana, 489
Texas:
  Dipodomys ordii, 743
  Onychomys leucogaster, 743
  Perognathus flavescens, 743
  Peromyscus (2 spp.), 743
  Reithrodontomys montanus, 743
Thermoenergetics:
  Antechinus stuartii, 81
Thermoregulation:
  Antechinus stuartii, 81
  bats from cloud forests, 445
  prairie voles and montane voles, 536
  Sciurus carolinensis, 467
Thomasomys:
  caudivarius, 834
  onkiro, 834
  silvestris, 834
Thomomys talpoides, 991
Thrichomys apereoides, 333, 345
Thylamys velutinus, 58
Ticks, parasites of small mammals in Mississippi, 1039
Tonatia, 49
Track tubes, 734
Trapping:
  bias, 728
  Blarina hylophaga, 280
  Peromyscus (2 spp.), 260
  Peromyscus leucopus, 728
  track tubes compared, 734
Trinidad:
  Saccopteryx bilineata, 526
Trophic guilds, 775
Trophy hunting, influence on mountain sheep, 682
Tuco-tuco, 843
Tundra:
  Ursus arctos, 188
Tursiops truncatus, 178
Ultrasonic calls:
  Microtus (2 spp.), 536
United States:
  Blarina brevicauda, 734
  Blarina carolinensis, 1039
  Blarina hylophaga, 280
  Canis (2 spp.), 754
  Canis latrans, 501
  Cervus elaphus nelsoni, 229
  Clethrionomys gapperi, 734
  Corynorhinus townsendii ingens, 590
  Dipodomys ordii, 743
  Enhydra lutris, 893
  Eumetopias jubatus, 973
  Glaucomys sabrinus, 734
  Marmota flaviventris, 576
  Martes americana, 734
  Microtus canicaudus, 947
  Microtus pinetorum, 1039
  Myotis sodalis, 585
  Napaeozapus insignis, 734
  Nasua narica, 934
  Neotoma floridana, 1039
  Neotoma fuscipes, 866
  Neotoma magister, 1058
  Neotoma mexicana, 20
  Ochrotomys nuttalli, 1039
  Odocoileus hemionus hemionus, 229
  Odocoileus virginianus, 31
  Onychomys leucogaster, 743
  Oreamonas americanus, 218
  Oryzomys palustris, 1039
  Perognathus flavescens, 743
  Peromyscus (2 spp.), 260, 614, 743, 1039
  Peromyscus leucopus, 267
  Peromyscus polionotus ammobates, 252
  Petrogale penicillata, 437
  Phoca vitulina, 1111
  Rattus norvegicus, 478, 1039
  Reithrodontomys humulis, 1039
  Reithrodontomys montanus, 743
  Sciurus carolinensis, 1039
  Sigmodon hispidus, 1039
  Sorex cinereus, 734
  Spermophilus lateralis, 1013
  Tadarida brasiliensis mexicana, 489
  Tamiasciurus hudsonicus, 734
  Tamias striatus, 734, 1039
  Thomomys talpoides, 991
  Ursus americanus, 1020
  Vulpes velox, 199
  Zapus hudsonius, 739
Urinary pheromones:
  Antechinus stuartii, 71
Ursus:
  americanus, 1020
  arctos, 188
  arctos yesoensis, 1026

Uruguay:
Oxymycterus (11 spp.), 408
Venezuela:
Anoura latidens, 445
Didelphis (3 spp.), 1087
Sturnira erythromos, 445
Tadarida brasiliensis, 445
Vocalizations:
Alouatta pigra, 159
Balantiopteryx (3 spp.), 1049
Carollia perspicillata, 516
dialect, 526
Microtus (2 spp.), 536
Saccopteryx bilineata, 526
Tamias striatus, 546
Voles:
Clethrionomys glareolus, 459
Ellobius talpinus, 459
Eothenomys (2 spp.), 852
Microtus (2 spp.), 536
Microtus canicaudus, 947
Microtus pinetorum, 1039
Vulpes:
velox, 199
zerda, 99
Wallaby:
Petrogale penicillata, 437
Washington:
Enhydra lutris, 893
Water influx:
in Oryx leucoryx, 665
Weather:
effect on horn growth in Capra ibex ibex, 245
White-eared opossum, 354
White-footed mouse, 267, 729, 1039
White-nosed coati, 934
White-tailed deer, 31, 884
Wolverine, 1141
Wolves:
dietary overlap with coyote, 754
Wood rat:
Allegheny, 1058
dusky-footed, 866
eastern, 1039
Wyoming:
Ursus americanus, 1020
Vulpes velox, 199
Yellow-bellied marmot, 576
Zacatuche rabbit, 1077
Zalophus californianus, 135
Zapus hudsonius, 734
Zimbabwe:
Canis (2 spp.), 599
Loxodonta africana, 652